Dermatology & Plastic Surgery SPECIALISTS

MICRO LASER PEEL

BEFORE YOUR TREATMENT

- Avoid sun exposure, tanning beds and self-tanner for at least **two weeks** prior to treatment.
- Discontinue retinol, tretinoin, adapalene and other exfoliating or irritating skincare products for **five to seven days prior**, or as directed by your provider.
- Avoid blood thinners (aspirin, ibuprofen, naproxen, fish oil, vitamin E, ginkgo, garlic, turmeric and other supplements) for **five to seven days prior**, if medically appropriate.
- Do not take alcohol for **twenty-four hours** before treatment.
- Arrive with clean skin. Do not wear makeup, skincare products, or lotion on the day of treatment.
- Inform your provider of all medications, supplements, allergies, or medical conditions.
- Alert your provider if you have a history of cold sores, keloid scarring or any skin conditions.
- Do not have waxing, chemical peels or other facial treatments for at least **two weeks** prior.
- Alert your provider if you have any neuromuscular or autoimmune disorders or known allergies.

Dermatology & Plastic Surgery SPECIALISTS

MICRO LASER PEEL

AFTER YOUR TREATMENT

- Expect redness, warmth and a mild sunburn-like sensation for **one to three days**.
- Skin may feel tight, dry and rough with light to moderate peeling for **three to seven days**.
- Keep skin clean and moisturized with a gentle, hydrating product as directed.
- Do not pick or peel flaking skin. Let it shed naturally.
- Avoid makeup for **twenty-four hours** (or until skin feels comfortable).
- Avoid retinol, exfoliants, acids or other active skincare products for **five to seven days**, or as directed by your provider.
- Avoid direct sun exposure, tanning beds and excessive heat for at least **two weeks**. Use a mineral sunscreen (SPF 30 or higher) daily.
- Avoid strenuous exercise, saunas, hot tubs and excessive heat for **twenty-four to forty-eight hours**.
- Use cool compresses and/or vinegar soaks as directed to help with discomfort and healing (see separate aftercare sheet).
- You may return to normal activities as tolerated.
- Contact us if you experience increased redness, swelling, blistering or any concerns.